



Department
for Environment
Food & Rural Affairs



FIELD GUIDE TO INVASIVE ALIEN PLANT PESTS IN THE CARIBBEAN UK OVERSEAS TERRITORIES



PART 1 – INTRODUCTION

Chris Malumphy, Sharon Reid, Rachel Down, Jackie Dunn
and Debbie Collins

FIELD GUIDE TO INVASIVE ALIEN PLANT PESTS IN THE CARIBBEAN UK OVERSEAS TERRITORIES

Part 1 Introduction

Chris Malumphy, Sharon Reid, Rachel Down, Jackie Dunn and Debbie Collins

Second Edition

Fera Science Ltd., National Agri-food Innovation Campus, Sand Hutton, York, YO41 1LZ, United Kingdom

<https://fera.co.uk/>

Published digitally: April 2018.

Second edition published digitally: April 2019.

Divided into 6 parts to enable easier download

© Crown copyright 2018-19

Suggested citation: Malumphy, C., Reid, S., Down, R., Dunn., J. & Collins, D. 2019. Field Guide to Invasive Alien Plant Pests in the Caribbean UK Overseas Territories. 2nd Edition. Part 1 – Introduction. Defra/Fera. 30 pp.

Frontispiece

Top row: Giant African land snail *Lissachatina fulica* © C. Malumphy; Mediterranean fruit fly *Ceratitidis capitata* © Crown copyright; Sri Lankan weevil, *Myloccerus undecimpustulatus undatus* adult © Gary R. McClellan. Second row: Cactus moth *Cactoblastis cactorum* caterpillar © C. Malumphy; Cottony cushion scale *Icerya purcashi* © Crown copyright; Red palm mite *Raoiella indica* adults © USDA. Third row: Tomato potato psyllid *Bactericera cockerelli* © Fera; Cotton bollworm *Helicoverpa armigera* © Crown copyright; Croton scale *Phalacroccoccus howertoni* © C. Malumphy. Bottom row: Red palm weevil *Rhynchophorus ferrugineus* © Fera; Tobacco whitefly *Bemisia tabaci* © Crown copyright; Brown marmorated stink bug *Halyomorpha halys* © David R. Lance, USDA APHIS PPO, Bugwood.org.

Contents

	Page
PART 1 – INTRODUCTION	
1. Purpose and scope	7
2. Introduction	7
2.1 UK Overseas Territories and biodiversity	7
2.2 Impact of invasive alien plant pests	9
2.3 Pathways of introduction	11
2.4 Mitigating plant health risks	12
2.5 Potential invasive alien invertebrate threats for each UKOT in the Caribbean	14
3. An introduction to invertebrate plant pests	16
3.1 Class Insecta – Insects	16
Order Coleoptera – Beetles	16
Order Diptera – True flies	19
Order Hemiptera – True bugs	20
Order Lepidoptera – Butterflies and moths	22
Order Thysanoptera – Thrips	25
Other groups of insects	26
3.2 Class Arachnida, Subclass Acari – Mites	27
3.3 Other groups of plant pests	28
Phylum Nematoda – Nematodes or round worms	28
Class Gastropoda – Slugs and snails	29
Class Diploda – Millipedes	30
PART 2 – PLANT DAMAGE	
4. Field diagnosis of plant damage caused by invertebrates	37
4.1 Damage to leaves	37
Leaves discoloured	37
Softer parts of the leaves eaten	42
Whole leaves or sections of leaves eaten or removed	43
Leaf mines	44
Abnormal growth	46
Leaf contamination	50
Premature leaf drop	54
4.2 Damage to fruit	54

Fruit discoloured	54
Fruit surface damaged	56
Abnormal growth	56
External contamination of fruit	57
Fruit with holes and bleeds	59
Internal tunnelling	60
4.3 Damage to trunk, branches and stems	61
Contamination of bark	61
Holes, sap flows and bleeds	63
Frass and sawdust on the bark or at the tree base	65
Tunnelling	66
4.4 Damage to whole tree and mortality	68
5. Specimen collection and preservation	69
5.1 Collecting methods	69
5.2 Preservation methods	70
PART 3 – INSECTS (except Hemiptera)	
6. Invasive Alien Invertebrate Species	79
INSECTA	
Coleoptera – Cerambycidae	
6.1 Brown mulberry longhorn beetle – <i>Apriona germari</i>	80
6.2 Monkey pod Round-headed Borer – <i>Xystrocera globosa</i>	83
Coleoptera – Curculionidae	
6.3 Polyphagous shot hole borer – <i>Euwallacea whitfordiodendrus</i>	86
6.4 Sri Lankan weevil – <i>Myloccerus undecimpustulatus undatus</i>	89
6.5 Red palm weevil – <i>Rhynchophorus ferrugineus</i>	92
6.6 Agave weevil – <i>Scyphophorus acupunctatus</i>	96
6.7 Mango seed weevil – <i>Sternochetus mangiferae</i>	100
Diptera – Drosophilidae	
6.8 Spotted wing Drosophila – <i>Drosophila suzukii</i>	103
Diptera – Tephritidae	
6.9 Fruitfly – <i>Bactrocera carambolae</i>	107
6.10 Oriental Fruitfly – <i>Bactrocera dorsalis</i>	111
6.11 Mediterranean Fruitfly – <i>Ceratitis capitata</i>	115
Hymenoptera – Formicidae	
6.12 Red imported fire ant – <i>Solenopsis invicta</i>	119

Isoptera – Rhinotermitidae	
6.13	Formosan subterranean termite – <i>Coptotermes formosanus</i> 123
Lepidoptera – Pyralidae	
6.14	Cactus moth – <i>Cactoblastis cactorum</i> 126
Lepidoptera – Noctuidae	
6.15	Cotton bollworm – <i>Helicoverpa armigera</i> 129
6.16	Fall armyworm – <i>Spodoptera frugiperda</i> 132
Thysanoptera – Phlaeothripidae	
6.17	White cedar thrips – <i>Holopothrips tabebuia</i> 136
Thysanoptera – Thripidae	
6.18	Western flower thrips – <i>Frankliniella occidentalis</i> 139
6.19	Melon thrips – <i>Thrips palmi</i> 143

PART 4 – HEMIPTERA (Bugs)

Hemiptera – Aleyrodidae	
6.20	Solanum whitefly – <i>Aleurotrachelus trachoides</i> 152
6.21	Tobacco whitefly – <i>Bemisia tabaci</i> 156
6.22	Cardin’s whitefly – <i>Metaleurodicus cardini</i> 159
6.23	Ficus whitefly – <i>Singhiella simplex</i> 162
Hemiptera – Aphididae	
6.24	Black citrus aphid – <i>Aphis (Toxoptera) citricidus</i> 165
6.25	Cypress aphid – <i>Cinara cupressi</i> 168
Hemiptera – Cixiidae	
6.26	American palm cixiid – <i>Haplaxius crudus</i> 172
Hemiptera – Liviidae	
6.27	Asian citrus psyllid – <i>Diaphorina citri</i> 176
Hemiptera – Pentatomidae	
6.28	Brown Marmorated Stink Bug – <i>Halyomorpha halys</i> 179
Hemiptera – Tingidae	
6.29	Avocado lace bug – <i>Pseudacysta perseae</i> 182
Hemiptera – Triozidae	
6.30	Tomato potato psyllid – <i>Bactericera cockerelli</i> 185

PART 5 – HEMIPTERA (Scale insects)

Hemiptera – Coccidae	
6.31	Pink wax scale – <i>Ceroplastes rubens</i> 195
6.32	Croton scale – <i>Phalacrocooccus howertoni</i> 198

6.33	Pine Tortoise Scale – <i>Toumeyella parvicornis</i>	201
	Hemiptera – Diaspididae	
6.34	Cycad Aulacaspis scale – <i>Aulacaspis yasumatsui</i>	205
6.35	Bermuda cedar scale or Minute cypress scale – <i>Carulaspis minima</i>	208
6.36	Cactus scale – <i>Diaspis echinocacti</i>	211
6.37	Lesser snow scale – <i>Pinnaspis strachani</i>	214
6.38	White Peach Scale – <i>Pseudaulacaspis pentagona</i>	217
	Hemiptera – Kerriidae	
6.39	Lobate lac scale – <i>Paratachardina pseudolobata</i>	221
	Hemiptera – Monophlebidae	
6.40	A fluted scale – <i>Crypticerya genistae</i>	224
6.41	Cottony cushion scale – <i>Icerya purchasi</i>	227
	Hemiptera – Ortheziidae	
6.42	Greenhouse orthezia or Kew bug – <i>Insignorthezia insignis</i>	231
6.43	Citrus ortheziid – <i>Praelongorthezia praelonga</i>	234
	Hemiptera – Pseudococcidae	
6.44	Pink hibiscus mealybug – <i>Maconellicoccus hirsutus</i>	237
6.45	Lantana mealybug – <i>Phenacoccus parvus</i>	241
6.46	Bougainvillea mealybug – <i>Phenacoccus peruvianus</i>	244
6.47	Solenopsis or cotton mealybug – <i>Phenacoccus solenopsis</i>	247

PART 6 INVERTEBRATES (Except insects) & REFERENCES

GASTROPODA

Stylommatophora – Achatinidae

6.48	Giant African Land Snail – <i>Lissachatina fulica</i>	257
------	---	-----

DIPLOPODA

Spirobolida – Rhinocricidae

6.49	Bumblebee millipede – <i>Anadenobolus monilicornis</i>	261
------	--	-----

ARACHNIDA

Trombidiformes – Tenuipalpidae

6.50	Red palm mite – <i>Raoiella indica</i>	264
------	--	-----

7.	Acknowledgments	267
8.	References	267
9.	Appendices	286
9.1	Major sources of further information	286
9.2	Fera Invertebrate plant pest identification Service for the UKOTs	286

1. Purpose and scope

Invasive alien species are a major threat to biodiversity globally and particularly to the United Kingdom Overseas Territories (UKOTs) in the Caribbean. Early detection and identification of invasive alien species can enable effective and appropriate measures to be taken in a timely manner to eradicate, contain and/or manage the pests. The purpose of this guide is to assist this process by:

- Raising awareness of the potential harm that invasive alien invertebrate plant pests can cause if accidentally introduced into the UKOTs in the Caribbean region;
- Providing pertinent information to enable those involved with plant health to recognise the main groups of invertebrate plant pests and the plant damage they cause;
- Providing detailed factsheets on key invasive alien invertebrate pests that currently threaten plant health in the region.

This guide is intended to be used by those working in biosecurity and plant health, and by all those interested in preserving the unique environments and biodiversity found in the Caribbean UKOTs. The guide will not enable the user to identify to species all the invasive invertebrate plant pests that they may encounter, but it does provide advice on how to collect and preserve specimens for further study and where to seek assistance.

2. Introduction

2.1 UK Overseas Territories and Biodiversity

There are 14 UKOTs spread across the World of which five are in the Caribbean (Anguilla, British Virgin Islands (BVI), Cayman Islands, Montserrat and the Turks and Caicos Islands (TCI)) and one located in the adjacent North Atlantic (Bermuda) (Fig. 2.1.1). The UKOTs support a diverse range of unique ecosystems and habitats (Fig. 2.1.2) and sustain many rare and threatened plant and animal species (Fig. 2.1.3). It is estimated that over 90% of the UK's biodiversity is found in its Overseas Territories, with more priority ecosystem types occurring in the UKOTs than in the whole of metropolitan UK (Defra, 2012).



Figure 2.1.1 Map showing the location of the UK Overseas Territories and Crown Dependencies © JNCC



Figure 2.1.2 Dry tropical scrub in Prickly Pear National Park, British Virgin Islands © C. Malumphy



Figure 2.1.3 An example of the unique fauna in the UKOTs, the Anegada ground iguana (*Cyclura pinguis*) is a critically endangered species only found in Anegada, British Virgin Islands © C. Malumphy

As the UKOTs are mostly small islands, their biodiversity has evolved in isolation, resulting in a high proportion of endemic species, found nowhere else in the world. To date, over 340 endemic species have been found in the UKOTs, compared to about 60 in metropolitan UK (Weighell & Pelembe, 2009). A review by Pienkowski (2005) noted 61 sites in the UKOTs that met the criteria for Ramsar designation, indicating the global significance of the UKOT wetland habitats.

Further information on the UKOTs in the Caribbean and Bermuda and their biodiversity can be obtained from the following links:

- Joint Nature Conservation Committee (JNCC) – <http://jncc.defra.gov.uk/page-4079>
- UK Overseas Territories Conservation Forum (UKOTCF) – <http://www.ukotcf.org/index.cfm>
- Policy UK Overseas Territories – <https://www.gov.uk/government/policies/uk-overseas-territories>
- Kew Royal Botanic Gardens – UK Overseas Territories Programme – <http://www1.kew.org/science/ukots/index.html>
- GB Non-Native Species Secretariat – UK Overseas Territories Home – <http://www.nonnativespecies.org/ots/otsMap.cfm>
- Government of Anguilla – <http://www.gov.ai/>
- Government of Bermuda – <https://www.gov.bm/>
- Cayman Islands Government – <http://www.gov.ky/portal/page/portal/cighome>
- Monserrat Government UK Office – montserrat-gov.org/
- Government of Turks and Caicos Islands – <https://www.gov.tc/>
- Government of the Virgin Islands – <http://www.bvi.gov.vg/>

2.2 Impact of invasive alien plant pests

Invasive alien species are a major cause of biodiversity loss globally. Their impacts are often most severe on island environments typical of the UKOTs. Island species or populations are often vulnerable to the impacts of non-native species because of their previous isolation from predators, diseases or competitors. Indeed, the majority (72%) of global extinctions since 1500 have occurred on islands and, for example, 67% of threatened bird species are found on oceanic islands. A review of non-native species in the UKOTs indicated that there are a minimum of 2261 non-native species recorded as occurring across the UK Overseas Territories and Crown Dependencies (JNCC, 2007). The UKOTs contain over 500 globally threatened species, and many undisturbed habitats of international significance. Invasive alien species not only impact biodiversity but can also have a negative economic and social impact as illustrated below.

Environmental impacts

The introduction of invasive alien invertebrate plant pests into the UKOTs can have an enormous negative impact on the environment, in some cases permanently changing the ecology. There have been a series of catastrophic declines of native plants in the UKOTs which are directly attributable to the introduction of invasive species, for example Bermuda's dominant tree species, Bermuda cedar (*Juniperus bermudiana*), declined by almost 95% between 1946 and 1956 following the introduction of Bermuda cedar scale (*Carulaspis minima*) (see Fact sheet 6.35) and to a lesser degree Newstead scale (*Lepidosaphes newsteadii*), despite various attempts at biological control (Challinor & Wingate, 1971; Wingate, 2001, 2011). Fortunately, some natural regeneration still occurs from the small numbers of remaining Bermuda cedar which exhibit some resistance to the scale insect. The Bermuda

cedar was a nesting habitat for birds and home to numerous insects and other invertebrates. Many species that were adapted to life in the cedar-dominated forest have seriously declined, such as the native bluebird (*Sialia sialis bermudensis*) and the endemic cicada (*Tibicen bermudiana*), which is now extinct. It is hoped that Bermuda cedar will survive long enough to develop an inherent resistance, as have the closely-related mainland junipers.

A similar catastrophic decline of about 90-95% of the dominant tree species, Caicos or Caribbean pine (*Pinus caribaea* var. *bahamensis*), has occurred in the Turks and Caicos Islands from about 2005 onwards, following the accidental introduction of pine tortoise scale (*Toumeyella parvicornis*) (Malumphy *et al.*, 2012) (Figs 2.2.1-2.2.2) (see Fact sheet 6.33). The Caicos pine is the foundation species of the pine yard habitat, a globally endangered ecosystem on which many other plants, birds, reptiles, and insects depend. Action is being taken through the Caicos Pine Recovery Project to safeguard the future of the Caicos pine.



Figure 2.2.1 Healthy *Pinus caribaea* forest in the Bahamas © C. Malumphy



Figure 2.2.2 Dead and dying *Pinus caribaea* trees in the Turks and Caicos Islands due to a massive infestation of tortoise scale *Toumeyella parvicornis* © Dr Martin Hamilton, Kew

More recently the endemic century plant (*Agave missionum*) has been almost wiped out from most areas of the British Virgin Islands following the introduction of the agave snout weevil (*Scyphophorus acupunctatus*) (see Fact sheet 6.6). This is currently being investigated by a team from Kew Gardens in collaboration with the National Parks Trust of the Virgin Islands.

Economic impacts

The Asian pink hibiscus mealybug, *Maconellicoccus hirsutus* (see Fact sheet 6.44), was first reported in the Caribbean in 1994 in Grenada. This was the first record of the insect as a major pest in the Americas. By the beginning of 2001, the pest had spread to over 25 territories from Guyana and Venezuela in the South to the Bahamas in the North (Williams, 1996). The pest has also extended its distribution to Central America and southern North America. It attacks more than 330 plant species (Chong, 2009), including many agricultural and horticultural crops. It caused estimated economic losses, which include crop losses, costs of control, and impact on trade, of \$18.3 million (USD) in Grenada (1995–1998), \$280,000 in Saint Kitts (1995–1997), \$5.1 million in Trinidad (1995–1997), \$67,000 in Saint Lucia and Saint Vincent, and \$3.4 million in the Grenadines (Kairo *et al.*, 2000). Early attempts to control the pest using pesticides resulted in failure and a classical biological control approach was adopted. Several exotic natural enemies were introduced but control was attributed to

the parasitoid wasp *Anagyrus kamali* and the ladybird *Cryptolaemus montrouzieri*. In all countries where biological control was implemented, this resulted in success in most areas.

More recent examples of invasive alien invertebrates in the Caribbean region are the papaya mealybug *Paracoccus marginatus*, which has become a pest in Antigua, Saint Kitts and Nevis, and the British and US Virgin Islands (Miller *et al.*, 1999); and the Asian citrus psyllid *Diaphorina citri* (see Fact sheet 6.27), which has become widespread in the Caribbean since 1998 (CABI/EPPO, 2011), and is a vector of the very serious citrus huanglongbing (greening) disease caused by the bacterium *Liberibacter asiaticus*. The latter disease has had a massive impact in Florida where commercial citrus production is no longer viable in many areas.

Social Impacts

The National tree of the Turks and Caicos Islands is the Caicos pine which has suffered a catastrophic decline following the introduction of pine tortoise scale (see section on environmental impact above).

The National tree of the British Virgin Islands is the White Cedar *Tabebuia heterophylla*. This tree was important because its wood was used to make the stern, stern posts and frame of the famous Virgin Islands sloops. The foliage of most white cedar plants has been severely galled by the recent introduction of the tabebuia thrips *Holopothrips tabebuia* (see Fact sheet 6.17) reducing their aesthetic appearance and even killing young plants.

The National tree of Bermuda is the Bermuda cedar which has suffered a catastrophic decline following the introduction of Bermuda cedar scale or minute cypress scale (see section on environmental impact above). The cedar was a valuable timber resource and of great cultural value to Bermudians.

The tradition of using a mature century plant *A. missionum* as a Christmas tree in the BVI, by spraying it with silver paint and decorating it, has disappeared as the plant has become rare following the introduction of the agave snout weevil *S. acupunctatus* (see Fact sheet 6.6).

There was a significant social (and economic) impact in many territories in the Caribbean following the introduction of the pink hibiscus mealybug, *M. hirsutus*, due to a reduction in agricultural production and exports (see Fact sheet 6.44).

2.3 Pathways of introduction

There is relatively little detailed quantitative data available on the long distance natural dispersal of invertebrates. However, it certainly occurs, being facilitated by migrating birds, mammals, and insects and by air and water currents (rafting). In the Caribbean hurricanes are likely to be important in distributing some arthropods, such as the red palm mite and pink hibiscus mealybug, over wide areas. However, natural pathways appear to be only a minor component of the recent and rapid global spread of invasive alien species in contrast with human-mediated dispersal. The main pathway for most plant pest species is plant trade, particularly the movement of plants for planting. The Caribbean is a major centre of tourism and receives large volumes of passengers and cargo by sea and air. Tourist developments require a regular supply of ornamental plants and as most of the UKOTs do not have a horticultural sector to meet local needs, there is a constant flow of ornamental plants into the territories. Florida is the major plant producer in the region and exports plants to all parts of the Caribbean. If an invasive pest is introduced into Florida, there is a potential pathway for it to spread rapidly throughout the region.

2.4 Mitigating plant health risks

A detailed discussion of mitigating plant health risks is beyond the scope of this guide. A simple outline of a plant health strategy is presented below to provide a pre- to post-border biosecurity continuum. The pre-border component is very important as it is always beneficial to reduce the likelihood of a pest being introduced in the first place rather than addressing the consequences after its introduction. It should also be stressed that biosecurity is the responsibility of everybody involved with plants including the agricultural, horticultural and tourist industries, landowners, non-governmental organizations and the public. There needs to be a continual awareness of the risks, vigilance and involvement by all the stakeholders for a biosecurity strategy to be effective.

Pre-Border

- Evidence based and proportionate plant health legislation (including regulated pests and plants) to ensure the best protection for plant health.
- International collaboration to reduce the likelihood of pests arriving at the border in the first place.
- Horizon scanning to identify potential new pests and gain advance warning and intelligence.
- Effective and prompt sharing of information on new risks and pathways.
- Careful sourcing and selection of plant material.

Border

- Border inspections to reduce the risk of pests crossing the border, and feedback intelligence to Pest Risk Assessors.
- Targeted risk-based inspections of imported plant material.
- Develop and validate new detection and identification methods.
- Enhance collaboration between border force and trade to gain intelligence of pathways.

Inland

- Targeted surveillance to enable early detection of new pests.
- Developing better ways to respond and reduce the chance of establishment.
- Develop contingency plans and clear governance to eradicate or successfully manage outbreaks.
- Development of more resilient horticultural and agricultural sectors to increase the ability to manage pests and reduce the need for imports.
- Building resilience in the UKOTs and learning to live with established pests.

Further information on biosecurity and plant health threats may be obtained online from the following links:

- **IPPC – International Plant Protection Convention**
<https://www.ippc.int/en/>

The IPPC is the phytosanitary standard setting organization referenced in the Sanitary and Phytosanitary Measures Agreement of the World Trade Organization (WTO). They have established International Standards for Phytosanitary Measures (ISPMs) (<https://www.ippc.int/en/core-activities/standards-setting/ispm/>) which provide guidance on all aspects of plant health. The IPPC also provides Phytosanitary Resources (<http://www.phytosanitary.info/>) including e-learning

modules, manuals, training materials, diagnostic protocols, videos, advocacy materials, photographs, a roster of consultants and databases of projects and activities.

- **North American Plant Protection Organization (NAPPO)**
<https://www.nappo.org/index.php>

NAPPO is the phytosanitary standard setting organization recognized by the North American Free Trade Agreement (NAFTA). It was created in 1976 as a regional organization of the IPPC. One of the many useful links on this site is the Phytosanitary Alert System (PAS) which provides up-to-date information on plant pest situations of significance to North America, most of which are relevant to the Caribbean UKOTs (<http://www.pestalert.org/index.cfm?NAPPOLanguagePref=%27English%27>).

- **Florida Department of Agriculture and Consumer Services**
<https://www.freshfromflorida.com/>

There is a useful series of fact sheets on plant health threats (pests and diseases) available on the 'Plant Industry Pest Alerts' page (<https://www.freshfromflorida.com/Divisions-Offices/Plant-Industry/Plant-Industry-Publications/Pest-Alerts>)

- **European and Mediterranean Plant Protection Organization (EPPO)**
<https://www.eppo.int/>

EPPO is an intergovernmental organization responsible for cooperation and harmonization in plant protection within the European and Mediterranean region. Under the International Plant Protection Convention (IPPC), EPPO is the regional plant protection organization (RPPO) for Europe.

2.5 Major invertebrate threats for each of the UKOTs in the Caribbean

A Horizon Scanning workshop was convened in the Cayman Islands during 2018 to identify the most important economic, human and biodiversity threats to the UKOTs in the Caribbean. Invertebrate species identified as major potential threats, currently absent, are listed below for each territory with the area of potential impact and the part and page number to the appropriate fact sheet.

Anguilla

Species	Impact	Part	Page
Fruitfly – <i>Bactrocera carambolae</i>	Economic	3	107
Mediterranean Fruitfly – <i>Ceratitis capitata</i>	Economic	3	115
Formosan subterranean termite – <i>Coptotermes formosanus</i>	Economic	3	123
Cactus moth – <i>Cactoblastis cactorum</i>	Biodiversity	3	126
Asian citrus psyllid – <i>Diaphorina citri</i>	Economic	4	176

Bermuda

Species	Impact	Part	Page
Sri Lankan weevil – <i>Myllocerus undecimpustulatus</i>	Biodiversity	3	89
Agave weevil – <i>Scyphophorus acupunctatus</i>	Biodiversity	3	96
Mediterranean Fruitfly – <i>Ceratitis capitata</i>	Economic	3	115
Red imported fire ant – <i>Solenopsis invicta</i>	Biodiversity	3	119
Formosan subterranean termite – <i>Coptotermes formosanus</i>	Economic	3	123
Asian citrus psyllid – <i>Diaphorina citri</i>	Economic	4	176
Croton scale – <i>Phalacrocooccus howertoni</i>	Biodiversity	5	198
Lobate lac scale – <i>Paratachardina pseudolobata</i>	Biodiversity	5	221
Pink hibiscus mealybug – <i>Maconellicoccus hirsutus</i>	Biodiversity	5	237
Giant African Land Snail – <i>Lissachatina fulica</i>	Biodiversity, Human Health, Economic	6	257
Red palm mite – <i>Raoiella indica</i>	Biodiversity, Economic	6	264

British Virgin Islands

Species	Impact	Part	Page
Fruitfly – <i>Bactrocera carambolae</i>	Economic	3	107
Mediterranean Fruitfly – <i>Ceratitis capitata</i>	Economic	3	115
Formosan subterranean termite – <i>Coptotermes formosanus</i>	Economic	3	123
Asian citrus psyllid – <i>Diaphorina citri</i>	Economic	4	176

Giant African Land Snail – <i>Lissachatina fulica</i>	Biodiversity, Human Health, Economic	6	257
---	--------------------------------------	---	-----

Cayman Islands

Species	Impact	Part	Page
Agave weevil – <i>Scyphophorus acupunctatus</i>	Biodiversity, Economic	3	96
Mango seed weevil – <i>Sternochetus mangiferae</i>	Economic	3	100
Fruitfly – <i>Bactrocera carambolae</i>	Economic	3	107
Mediterranean Fruitfly – <i>Ceratitis capitata</i>	Economic	3	115
Formosan subterranean termite – <i>Coptotermes formosanus</i>	Economic	3	123
Giant African Land Snail – <i>Lissachatina fulica</i>	Biodiversity, Human Health, Economic	6	257

Monserrat

Species	Impact	Part	Page
Agave weevil – <i>Scyphophorus acupunctatus</i>	Biodiversity	3	96
Fruitfly – <i>Bactrocera carambolae</i>	Economic	3	107
Mediterranean Fruitfly – <i>Ceratitis capitata</i>	Economic	3	115
Formosan subterranean termite – <i>Coptotermes formosanus</i>	Economic	3	123
Croton scale – <i>Phalacrocooccus howertoni</i>	Biodiversity	5	198
Lobate lac scale – <i>Paratachardina pseudolobata</i>	Biodiversity	5	221
Giant African Land Snail – <i>Lissachatina fulica</i>	Biodiversity, Human Health, Economic	6	257

Turks and Caicos Islands

Species	Impact	Part	Page
Agave weevil – <i>Scyphophorus acupunctatus</i>	Biodiversity	3	96
Mediterranean Fruitfly – <i>Ceratitis capitata</i>	Economic	3	115
Formosan subterranean termite – <i>Coptotermes formosanus</i>	Economic	3	123
Asian citrus psyllid – <i>Diaphorina citri</i>	Economic	4	176
Giant African Land Snail – <i>Lissachatina fulica</i>	Biodiversity, Human Health, Economic	6	257

3. An introduction to invertebrate plant pests

What is a plant pest? It is defined by the International Plant Protection Convention (IPPC) as ‘Any species, strain or biotype of plant, animal or pathogenic agent injurious to plants or plant products’ (IPPC, 2007). With this broad definition, any invertebrate that feeds on, or otherwise damages, any living or dead plant material is a plant pest.

What is an invertebrate? It is an animal that neither possess’ nor develops a vertebral column (commonly known as a backbone or spine). Familiar examples of terrestrial invertebrates include worms, slugs, snails, woodlice, land crabs, millipedes, centipedes, spiders, mites and insects. Within the UKOTs, the most important group of invertebrate plant pests are the insects, followed by the mites.

Accurate identification is important for biosecurity, pest management and risk assessment, as the correct name will link up with all the knowledge on the life cycle, host range, impact, natural enemies and appropriate management practices. An incorrect identification can result in the wrong pest risk assessment and inappropriate management action which can exacerbate the situation. The identification is particularly critical if a biological control program is going to be implemented.

3.1 Class Insecta – Insects

The great majority of insect plant pests are assigned to just five orders of insects: Coleoptera, Diptera, Hemiptera, Lepidoptera, and Thysanoptera. If you can recognize these five groups, you will be able to place the majority of insect plant pests that you encounter in the field into the correct Order. The main characters used to distinguish these orders are summarised in Table 3.1.

Order Coleoptera – Beetles

Figs 3.1.1-3.1.4.

This is the largest and most diverse order in the entire animal kingdom, with about 370 thousand species worldwide. Beetles are minute to large insects, normally with two pairs of wings of which the front ones (called elytra) are hard or leathery and meet neatly along the mid-line of the dorsal surface. The hind wings are membranous and folded out of sight beneath the elytra. The prothorax (section behind the head) is normally large and mobile. The mouthparts are of the biting type. Beetles are usually instantly recognizable although some bugs (Hemiptera: Heteroptera) are similar in appearance but may be distinguished by their overlapping front wings and sucking mouthparts. Beetles have a complete life-cycle with eggs, larvae (grubs), pupa and adults. This is the largest order of insects and contains many highly invasive plant pests.

Examples of beetle families that contain major plant pests include:

Family **Bostrychidae** – Auger beetles, False powder-post beetles, or Horned powder-post beetles
This family contains several pantropical wood boring and storage pests.

Family **Bruchidae** – Seed beetles

Many bruchids attack stored seeds such as beans and lentils.






<p>Coleoptera – Beetles</p> <p>Minute to very large insects, usually with hardened bodies. Forewings characteristically modified into hardened cases (elytra), which fold over and protect the membranous hindwings when at rest. Mouthparts are adapted for chewing. The larvae are known as grubs.</p>	
<p>Diptera – True flies</p> <p>Generally small insects, easily recognized by the membranous forewings in combination with hind wings that are modified to form a pair of small club-shaped halteres used for balancing when flying. Mouthparts are adapted for sucking. The larvae are known as maggots.</p>	
<p>Hemiptera – True bugs</p> <p>Small to medium sized insects with sucking mouthparts, two pairs of wings or reduced hind wings. In the Auchenorrhyncha and Sternorrhyncha, both pairs of wings (if present) are membranous; in the Heteroptera, the bottom half of the forewings is thickened, forming the hemelytra. The immature stages are known as nymphs.</p>	
<p>Lepidoptera – Butterflies and moths</p> <p>Minute to very large insects with elongate antennae and two pairs of large membranous wings, the entire body and wings are covered by scales. Mouthparts are generally elongated to form a long sucking proboscis, held coiled beneath the head when not in use. Many butterflies and moths are brightly coloured. The larvae are known as caterpillars.</p>	
<p>Thysanoptera – Thrips</p> <p>Minute insects with slender bodies and, usually two pairs of narrow, fringed wings. Sucking mouthparts. Commonly found in flowers. The immature stages are known as larvae.</p>	

Table 3.1 Key characters used to recognize the five main orders of insect that contain the majority of invasive alien plant pests © Crown copyright.

Family **Buprestidae** – Jewel beetles (Fig. 3.1.2)

This is a large, mainly tropical family, containing some beautiful brightly metallic coloured insects. The larvae develop in wood and some species are major invasive forestry pests.

Family **Cerambycidae** – Longhorn beetles (Fig. 3.1.1)

There are over 26 thousand described species, and this is one of the most economically important families of beetles as the larvae tunnel into wood. Some are major pests of timber and forestry. They are usually recognizable by their extremely long antennae and the family contains some of the largest of all known insects.

Family **Chrysomelidae** – Leaf beetles (Fig. 3.1.3)

This is a large family containing more than 37 thousand species of leaf-feeding beetle, most of which are quite small. It contains many important invasive plant pests.

Family **Curculionidae** – Weevils, bark beetles, ambrosia beetles (Fig. 3.1.4)

This is a huge family with more than 82 thousand species. Most can be recognized by the well-marked rostrum or snout and elbowed, clubbed antennae. Many of the species are covered in tiny, brightly-coloured scales. The bark beetles (Scolytinae) are typically cylindrical and have a reduced rostrum. Ambrosia beetles carry a fungus on which they feed. The larvae of most species feed enclosed in roots, stems, seeds or under bark. Many species are serious pests of trees and stored products.

Family **Scarabaeidae** – Chafers, dung beetles, scarabs

This is one of the largest of all insect families, with over 19 thousand species. It also contains some of the largest and heaviest insects, and many species have beautiful metallic colouration, and bear extraordinary 'horns' that develop on the head and thorax.



Figure 3.1.1 Red-necked longhorn beetle *Aromia bungii* (Cerambycidae) on *Rosa*, Italy. The very long antennae are characteristic for the family © C. Malumphy



Figure 3.1.2 Jewel beetle *Ctenoderus maulicus* (Buprestidae), Falkland Islands © Crown copyright



Figure 3.1.3 Adult *Eurypepla jamaicensis* (Chrysomelidae), Cayman Islands © C. Malumphy



Figure 3.1.4 *Diaprepes abbreviatus* (Curculionidae) on *Guaiacum officinale*, Cayman Islands © C. Malumphy

Order Diptera – True flies

Figs 3.1.5-3.1.8.

True flies are minute to medium sized insects with a single pair of wings (forewings) used to fly. The hindwings having evolved into minute dumbbell-shaped organs known as halteres, which act as high-speed sensors of rotational movement and allow dipterans to perform advanced aerobatics. Diptera is a large order with about 125 thousand species described. They have sucking mouthparts and a complete life-cycle with eggs, larvae (commonly known as maggots), puparia and adults.

Examples of fly families that contain major plant pests include:

Family **Agromyzidae** – Leaf mining flies (Fig. 3.1.5)

There are approximately 2,500 species and the larvae mine the foliage of various plants.

Family **Cecidomyiidae** – Gall midges (Fig. 3.1.6)

There are approximately six thousand species and the larvae of most gall midges feed within plant tissue, creating abnormal plant growths called galls.



Figure 3.1.5 Chrysanthemum leafminer *Chromatomyia syngenesiae* (Agromyzidae), Falkland Islands © C. Malumphy



Figure 3.1.6 Olive midge *Lasioptera berlesiana* (Cecidomyiidae) reared from a gall on olive *Olea europea*, Lebanon © Crown copyright



Figure 3.1.7 Vinigar fly *Zaprionus indianus* (Drosophilidae) reared from *Punica*, India, invasive in Europe © Crown copyright



Figure 3.1.8 West Indian fruitfly *Anastrepha obliqua* (Tephritidae) reared from *Mangifera indica*, Dominican Republic © Crown copyright

Family **Tephritidae** – Fruit flies (Fig. 3.1.8)

There are nearly five thousand described species of fruit fly. They are small to medium-sized flies that are often colourful, and usually have pictured or patterned wings. This is one of the most economically important groups of insects in the tropics and subtropics because the larvae of most species develop in plant tissue, especially fruit. Many species are highly invasive.

Order **Hemiptera** – True bugs

Figs 3.1.9-3.1.16

The Hemiptera comprise some 50 to 80 thousand species and are one of the most morphologically and biologically diverse orders of insects. Most hemipterans feed on plants, using their sucking and piercing mouthparts to extract plant sap. Some are parasitic while others are predators that feed on other insects or small invertebrates. They have an incomplete life-cycle with eggs, nymphs and adults. One exception is the whitefly (Aleyrodidae) where the final (4th) nymphal instar is called a pupa or puparium. Whitefly taxonomy is based on the morphology of this stage rather than the adult which is used for the rest of the Hemiptera. Many hemipterans vector plant pathogenic pathogens and many egest honeydew which serves as a medium for the growth of black sooty moulds. Hemipterans are very common in the UKOTs in the Caribbean.

Examples of hemipteran families that contain plant pests include:

Sub-order Heteroptera

Family **Pentatomidae** – Shield bugs or stink bugs (Fig. 3.1.9)

Large family of often brightly coloured, medium sized to large bugs with shield-shaped bodies. All are armed with stink glands.

Family **Tingidae** – Lace bugs (Fig. 3.1.10)

Most species are easily recognized by the delicate lace-like sculpturing of their wings and thorax. Most species are host specific.

Sub-orders Auchenorrhyncha and Sternorrhyncha

Family **Aleyrodidae** – Whiteflies (Fig. 3.1.11)

Very small insects with their body and wings covered in fine powdery white wax. Some species have banded wings or completely dark wings. The nymphs resemble scale insects (see below). Some species are important vectors of plant pathogenic viruses.

Family **Aphididae** – Aphids (Fig. 3.1.12)

Small, pear-shaped bugs. Wings (when present) are usually clear. Many species have two tubes (siphunculi) sticking out of their abdomen. They often have complicated life cycles with sexual and parthenogenetic generations, winged and wingless forms and alternating hosts. Some species are important vectors of plant pathogenic viruses.

Family **Cicadellidae** – Leaf hoppers or sharp shooters (Fig. 3.1.15)

Large family of mainly small bugs, many brightly coloured. This is one of the most economically important families due to the ability of many species to vector plant pathogens.

Family **Flatidae** – Plant hoppers or Moth bugs (Fig. 3.1.16)

Usually easily identified by very broad wings, often greenish but may appear grey due to a covering of wax. The nymphs often carry conspicuous tufts of white wax coming from the tip of their abdomen.

Families **Aphalaridae**, **Psyllidae**, **Triozidae** and others – Plant suckers or jumping plant lice

Small to minute bugs with clear wings, that look like miniature cicadas. Many are host specific and feed on woody plants. The adults jump.

Super-family **Coccoidea** – Scale insects

Small to minute plant parasitic bugs assigned to several families. Adult females are wingless, some are mobile (Ensign scale insects and mealybugs) but most are immobile in the adult female stage (soft and armoured scales). Many produce protective waxy covers. Scale insects are very common in the tropics.

Family **Asterolecaniidae** – Pit scalesFamily **Coccidae** – Soft scale insects (Fig. 3.1.14)Family **Diaspididae** – Armoured scale insectsFamily **Monophlebidae** – Giant scale insects (Fig. 3.1.13)Family **Ortheziidae** – Ensign scale insectsFamily **Pseudococcidae** – Mealybugs

Figure 3.1.9 Southern green shieldbug *Nezara viridula* (Pentatomidae), Greece © C. Malumphy



Figure 3.1.10 Adult lace bug *Phymacysta tumida*, Cayman Islands © C. Malumphy



Figure 3.1.11 Giant whitefly *Aleyrodicus* sp. (Aleyrodidae), BVI © C. Malumphy



Figure 3.1.12 Oleander aphid *Aphis nerii* (Aphididae), Bahamas © C. Malumphy



Figure 3.1.13 A giant scale *Crypticerya genistae* (Monophlebidae), BVI © C. Malumphy



Figure 3.1.14 Barnacle scale *Ceroplastes cirripediformis* (Coccidae), BVI © C. Malumphy



Figure 3.1.15 *Apogonalia histrio* adult on *Jatropha* (Cicadellidae), Cayman Islands © C. Malumphy



Figure 3.1.16 Sea grape flatid *Petrusa epilepsis* (Flatidae), BVI © C. Malumphy

Order Lepidoptera – Butterflies and moths

Figs 3.1.17-3.1.26.

There are about 180 thousand species assigned to the order Lepidoptera in 126 families, which account for 10% of the total described species of living organisms. It is one of the most widespread and instantly recognizable insect orders in the world, commonly known as the butterflies and moths. Adult lepidopteran species are characterized by several features, including scales (flattened hairs)

covering their bodies and wings, and a proboscis (sucking mouthparts). The scales give butterflies and moths their extraordinary variety of colours and patterns. Almost all species have some form of membranous wings, except for a few that have reduced wings or are wingless. They have a complete lifecycle with eggs, caterpillars (or 'worms' in the Caribbean), pupae (or chrysalis if a butterfly), and the adult. Mating and the laying of eggs are carried out by adults, normally near or on the larval host plant. The caterpillars are completely different from their adult forms, having a cylindrical body with a well-developed head, mandible (biting) mouth parts, three pairs of thoracic legs and from none up to five pairs of prolegs. As they grow, these larvae change in appearance, going through a series of stages called instars. Once fully matured, the larva develops into a pupa. A few butterflies and many moth species spin a silk case or cocoon prior to pupating, while others do not, instead going underground. Once the pupa has completed its metamorphosis, a sexually mature adult emerges.

Examples of lepidopteran families that contain major plant pests include:

Family Crambidae – Grass moths (Fig. 3.1.22)

Small moths which are variable in appearance. Some rest with closely folded wings and are inconspicuous, while others are brightly coloured and patterned, and rest with their wings spread. This family is closely related to the Pyralidae (see below) and many species are pests of crops while others are beneficial, helping to control invasive weeds.

Family Gracillariidae – Leaf mining moths

Small moths whose larvae mine leaves and stems. It contains several economic, horticultural and environmental pests. Some species are highly invasive.

Family Noctuidae – Owlet moths, cutworms or armyworms (Fig. 3.1.24)

This is a large family containing more than 11 thousand species. The adults are medium sized to large moths and are typically drab or camouflaged in colour. Many species are considered major agricultural pests around the world. Their larvae are typically known as "cutworms" or "armyworms" due to enormous swarms that destroy crops, orchards and gardens every year.

Family Pyralidae – Pyralid or snout moths (Fig. 3.1.26)

The adults are medium sized to large moths. Many species are economically important agricultural and storage pests.

Family Sesiidae – Clearwing moths (Fig. 3.1.21)

The adults of most sesiids have parts of their wings with hardly any of the normal lepidopteran scales, leaving them transparent. Many have bodies generally striped with yellow, red or white and they resemble wasps or hornets. Sesiid larvae typically bore into wood or burrow in plant roots and many are serious pests of fruit-tree or timber cultivation.

Family Sphingidae – Hawk moths, sphinx moths or hornworms (Fig. 3.1.25)

The adults are moderate to very large in size and are distinguished among moths for their rapid, sustained flying ability. Their narrow wings and streamlined abdomens are adaptations for rapid flight. The caterpillars are equally large and impressive. Most only feed on a narrow range of hosts and they are only occasional pests.

Family Tortricidae – Tortrix moths or leafroller moths

This is a large family containing more than 10 thousand species. They are small or medium sized moths that rest with their wings folded back, producing a rather rounded profile. Many of these are economically important agricultural pests.



Figure 3.1.17 Zebra longwing butterfly *Heliconius charithonia* (Nymphalidae), BVI © C. Malumphy



Figure 3.1.18 Gulf fritillary *Agraulis vanillae insularia* (Nymphalidae), BVI © C. Malumphy



Figure 3.1.19 Pale swallowtail butterfly *Papilio eurymedon* (Papilionidae), USA © C. Malumphy



Figure 3.1.20 Small tortoiseshell butterfly *Aglais urticae* (Nymphalidae), UK © C. Malumphy



Figure 3.1.21 Texas wasp moth *Horama panthalon* (Sesiidae), attracted to a light, BVI © C. Malumphy



Figure 3.1.22 Crambid moth *Samea ecclesialis* (Crambidae), attracted to a light, BVI © C. Malumphy



Figure 3.1.23 *Empyreuma pugione* (Erebidae) on *Ixora* flower, Cayman Islands © C. Malumphy



Figure 3.1.24 Army worm *Spodoptera* sp. (Noctuidae) caterpillar, BVI © C. Malumphy



Figure 3.1.25 Tetrio sphinx *Pseudosphinx tetrio* (Sphingidae) caterpillar, Saint Lucia © C. Malumphy



Figure 3.1.26 Cactus moth *Cactoblastis cactorum* (Pyralidae) caterpillar, BVI © C. Malumphy

Order Thysanoptera – Thrips

Figs 3.1.27-3.1.28

Thrips are minute, slender insects with fringed wings and unique asymmetrical mouthparts. Most species feed on plants by puncturing and sucking up the contents, although a few are predatory. Approximately six thousand species have been described. They are weak fliers but are frequently moved with plant trade. Many species are pests of commercially important crops. A few species serve as vectors for plant pathogenic viruses, especially the Tospoviruses. Some species of thrips are beneficial as pollinators or as predators of other insects or mites. In the right conditions, such as in greenhouses, many species can exponentially increase in population size and form large swarms because of a lack of natural predators coupled with their ability to reproduce asexually. Identification to species is usually only possible morphologically by examination of a slide-mounted specimen under a high-power microscope by a specialist.

Family **Phlaeothripidae** – thrips (Fig. 3.1.27)

This family contains approximately 3400 species and may be distinguished from the Thripidae by the last abdominal segment being modified into a tube-like structure, and generally being a larger size. Many species are harmless fungi feeders, but many others induce plant galls in which they complete their life-cycle.

Family Thripidae – thrips (Fig. 3.1.28)

This family contains more than 2000 species and may be distinguished from other thrips by a saw-like ovipositor curving downwards, narrow wings with two veins, and antennae of six to ten segments with stiletto-like forked sense cones on antennal segments III and IV. They generally live and feed within narrow spaces at the base of leaves and within flowers. Several species are major economic agricultural pests.



Figure 3.1.27 Cuban laurel thrips *Gynaikothrips ficorum* (Phlaeothripidae) adult female and eggs inside a leaf gall on *Ficus microcarpa*, Cyprus © C. Malumphy



Figure 3.1.28 Greenhouse thrips *Heliothrips haemorrhoidalis* (Thripidae) adult on *Laurus nobilis*, UK © C. Malumphy

Other groups of insects

Order Hymenoptera – Ants, bees, wasps and sawflies

Figs 3.1.29-3.1.30

Family Formicidae – Ants (Figs 3.1.29-3.1.30)

This family contains more than 12 thousand species. They are easily identified by their elbowed antennae and the distinctive node-like structure that forms their slender waists. They are eusocial insects that divide labour among castes consisting of sterile female ‘workers’, ‘soldiers’ and other specialised groups. Nearly all ant colonies also have one or more fertile females called ‘queens’ and some fertile males called ‘drones’. The majority of ants are not direct plant pests, although some, such as leaf cutter ants, can be important agricultural pests. Many species are highly invasive and invade buildings.

Order Isoptera – Termites or white ants

Figs 3.1.31-3.1.33

There are more than three thousand species of termite described. They are eusocial insects that divide labour among castes consisting of sterile male and female ‘workers’ and ‘soldiers’. All colonies have fertile males called ‘kings’ and one or more fertile females called ‘queens’. They usually feed on dead plant material and cellulose, generally in the form of wood, leaf litter, soil or animal dung. Most species are not plant pests and are important in recycling of wood and plant matter. Many species are economically significant as they can cause serious damage to buildings, crops, or plantation forests. They have an incomplete life-cycle and biting mouthparts. Their soil trails or mud shelter tubes and nests are ubiquitous in the Caribbean.



Figure 3.1.29 Golden carpenter ant *Camponotus sericeiventris* (Formicidae), Brazil © Crown copyright



Figure 3.1.30 Acrobat ants *Crematogaster scutellaris* (Formicidae), Italy © Crown copyright



Figure 3.1.31 Termite nest in tree, Mexico © C. Malumphy



Figure 3.1.32 Termite soil trails or mud shelter tubes on a tree, Mexico © C. Malumphy



Figure 3.1.33 Termite nest in tree, BVI © C. Malumphy



Figure 3.1.34 Brown headed bird locust *Acanthacris ruficornis* (Acrididae), Kenya © Fera



Figure 3.1.35 Painted grasshopper *Poekilocerus pictus* (Pyrgomophidae), India © Crown copyright

Order Orthoptera – Grasshoppers, locusts, crickets and Katydid

Figs 3.1.34-3.1.35

There are more than 20 thousand species of Orthoptera described worldwide. They have an incomplete metamorphosis, biting mouthparts and produce sound by rubbing (stridulating) various

parts of their wings and/or legs against each other. They cannot fold their wings, which are held along their bodies, and most have enlarged hind legs for jumping. Some locusts form large swarms and are important agricultural and horticultural pests.

3.2 Class Arachnida, Subclass Acari – Mites

Figs 3.2.1-3.2.2

The subclass Acari (or Acarina) is a diverse group of arachnids that contains mites and ticks. Adult mites are very small, most have four pairs of legs and their mouthparts may be adapted for biting, stinging, sawing or sucking. Mites occupy a wide range of ecological niches and feed on a wide variety of materials including living and dead animal, plants and fungi and, lichens and some are parasites of plants and animals. There are more than 50 thousand species described worldwide.

Family Eriophyidae – Gall or rust mites

This is a large family containing about 3,600 species of plant parasites, many induce plant galls. They are dispersed naturally by wind. Many are major agricultural pests.

Family Tenuipalpidae – Flat mites or False spider mites

This family is closely related to the spider mites and contains several major agricultural and horticultural pests.

Family Tetranychidae – Spider mites (Figs 3.2.1-3.2.2)

There are over a thousand species of spider mite. They generally feed on the lower surface of leaves where they spin protective silk webs. Many species of spider mites are major agricultural and horticultural pests.



Figure 3.2.1 *Tetranychus* sp. (Tetranychidae) on a shrub, BVI © C. Malumphy



Figure 3.2.2 Lewis mite *Eotetranychus lewisi* (Tetranychidae) feeding damage on *Euphorbia*, UK © Fera

3.3 Other groups of invertebrate plant pests

Phylum Nematoda – Nematodes or round worms

Figs 3.3.1-3.3.2

Nematodes are microscopic worms that shed their cuticle as they grow. They are one of the most important groups of invertebrate plant pests as many species parasitize plants. They are often overlooked as they are microscopic, difficult to detect and identify, and the symptoms they cause are

often non-specific. Field identification of nematodes is beyond the scope of this guide and the following paragraph merely provides a brief introduction to the group.

Nematodes feed on all parts of the plant, including the roots, stems, leaves, flowers and seeds. They feed in a variety of ways and may be endo- or ectoparasitic but all use a specialized spear-like mouthpart called a stylet. Nematodes often pierce and withdraw the contents of plant cells, killing them in the process. When this type of feeding occurs, large lesions are formed in the plant tissue. Some nematodes do not kill the plant cells they feed upon but induce the plant cells to enlarge and grow, thus producing one or more nutrient-rich feeding cells (giant cells) for the nematode. Many plant-parasitic nematodes feed on the roots of plants. The feeding process damages the plant's root system and reduces the plant's ability to absorb water and nutrients. Typical nematode damage symptoms are a reduction of root mass, a distortion of root structure and/or enlargement of the roots. Nematode damage of the plant's root system also provides an opportunity for other plant pathogens to invade the root and thus further weaken the plant. Direct damage to plant tissues by shoot-feeding nematodes includes reduced vigour, distortion of plant parts, and death of infected tissues depending upon the nematode species. The aboveground symptoms of nematode damage to roots are relatively generic or nondescript, including nutrient deficiency, incipient wilt, stunting, poor yield and sometimes plant death. Few diagnostic signs and symptoms of plant damage by nematodes exist except root galls, cysts, 'nematode wool', and seed galls. Thus, damage to crops by root-infesting nematodes often goes unnoticed by growers. Field patterns of nematode damage to roots begin in a small area and spreads radially from the initial infection site, often assisted by farm equipment. The only way to accurately diagnose nematode disease is to sample soil and plant material from suspected sites, extract nematodes for analysis and examine under a microscope.




Figure 3.3.1 Leaf and bud nematode *Aphelenchoides ritzemabosi* feeding damage on *Weigela*, UK © Crown copyright




Figure 3.3.2 Leaf and bud nematodes *Aphelenchoides* sp. in solution, extracted from leaf tissue, UK © Crown copyright

Class Gastropoda – Slugs and snails

Figs 3.3.3-3.3.6

Gastropods are found in marine and fresh water environments and on land. Terrestrial slugs and snails are frequently moved with plant trade and some species are invasive. Snails can be readily recognized as they have a hard-coiled shell in which they can completely retract their bodies for protection. Slugs do not possess an external shell. Only a relatively small number of slug and snail species are agricultural and horticultural pests, for example the Cuban brown snail (Fig. 3.3.3).



Figure 3.3.3 Cuban brown snail *Zachrysis provisoria* (Camaenidae), invasive pest in many Caribbean Islands, Cayman Islands © C. Malumphy



Figure 3.3.4 *Neritina (Vitta) virginea* (Neritidae), TCI © Crown copyright



Figure 3.3.5 Unidentified snail, BVI © C. Malumphy



Figure 3.3.6 Leatherleaf slug *Veronicella* sp. (Veronicellidae), Cayman Islands © C. Malumphy

Class Diplopoda – Millipedes

Figs 3.3.7-3.3.8

Millipedes are arthropods that are characterised by having two pairs of jointed legs on most body segments. Most millipedes have elongated cylindrical or flattened bodies with more than 20 body segments, while pill millipedes are shorter and can roll into a ball. Most millipedes are detritivores, that is they eat decaying leaves and other dead plant matter, and are thus not plant pests. Some can cause severe damage to seedlings while others can occur in huge numbers after rain and invade buildings. They can therefore be plant pests or cause a nuisance.



Figure 3.3.7 Giant black millipede *Anadenobolus arboreus* (Rhinocricidae), BVI © C. Malumphy



Figure 3.3.8 Rusty millipede *Trigoniulus corallinus* (Trigoniulidae), Cayman Islands © C. Malumphy